

**Synthesis, and Characterization and In-Silico Analysis Against SARS CoV-2
of Novel Benzimidazolium Salts**

Elvan ÜSTÜN^{1*}, Neslihan ŞAHİN²

¹Department of Chemistry, Faculty of Art and Science, Ordu University, 52200, Ordu, Turkey

²Department of Science Education, Faculty of Education, Cumhuriyet University, 58140, Sivas, Turkey.

*Corresponding Author: elvanustun77@gmail.com

Supplementary Files

Supplementary Files

Content	Page
Figure S1. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrums of 1	3
Figure S2. FT-IR spectrum of 1	4
Figure S3. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrums of 2	5
Figure S4. FT-IR spectrum of 2	6
Figure S5. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrums of 3	7
Figure S6. FT-IR spectrum of 3	8





Figure S1. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of **1**




Figure S2. FT-IR spectrum of **1**




Figure S3. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrums of **2**




Figure S4. FT-IR spectrum of **2**





Figure S5. ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrums of **3**




Figure S6. FT-IR spectrum of **3**